

Recommended Labs for PrEP Using F/TDF or F/TAF

Test	Initiation	Q3 months	Q6 months	As Indicated
HIV-Ab/Ag	<input type="checkbox"/>	<input type="checkbox"/>		
HIV-RNA	If ANY of the following: <input type="checkbox"/> Used PO PrEP in past 3mo <input type="checkbox"/> Used IM PrEP in past 12mo <input type="checkbox"/> Concern for HIV infection during ITRI ¹			<input type="checkbox"/> Concern for HIV infection during ITRI <input type="checkbox"/> At discontinuation
Complete Metabolic Panel	<input type="checkbox"/>		<input type="checkbox"/>	<input type="checkbox"/> eCrCl concerns <input type="checkbox"/> ALT concerns <input type="checkbox"/> At discontinuation
STI Panel ²	<input type="checkbox"/>	<input type="checkbox"/>		<input type="checkbox"/> For test-of-cure or monitoring <input type="checkbox"/> Each pregnancy <input type="checkbox"/> By patient request <input type="checkbox"/> At discontinuation
Pregnancy Test	<input type="checkbox"/>	<input type="checkbox"/>		<input type="checkbox"/> Concern for pregnancy & patient request <input type="checkbox"/> At discontinuation
Hepatitis A Ab Total	<input type="checkbox"/>			
Hepatitis B Panel ³	<input type="checkbox"/>			<input type="checkbox"/> HBsAg w/ each pregnancy
Hepatitis C Ab ⁴	<input type="checkbox"/>			<input type="checkbox"/> More frequent for PWIDs <input type="checkbox"/> Each pregnancy <input type="checkbox"/> Persistent ALT > 23IU/L

Footnotes

1. ITRI = Inter-test reactivity interval. The time between HIV-RNA positivity and HIV-Ag/Ab positivity
2. Syphilis (reverse screening algorithm).
Chlamydia and Gonorrhea at all applicable sites (genitourinary / oropharyngeal / rectal).
3. Surface antigen (HBsAg) + surface antibody (HBsAb) + core antibody (HBcAb)
4. If HCV-Ab is positive, obtain HCV-RNA.