

## Recommended Labs for PrEP Using Injectable Cabotegravir

Test	Initiation	At 1mo f/u	Q2 Months	As Indicated
HIV-RNA	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/> Concern for HIV infection <input type="checkbox"/> <b>At discontinuation</b>
Complete Metabolic Panel	<input type="checkbox"/>			<input type="checkbox"/> ALT concerns
STI Panel <sup>1</sup>	<input type="checkbox"/>			<input type="checkbox"/> For test-of-cure or monitoring <input type="checkbox"/> Each pregnancy <input type="checkbox"/> By patient request <input type="checkbox"/> <b>At discontinuation</b>
Pregnancy Test	<input type="checkbox"/>			<input type="checkbox"/> Concern for pregnancy & patient request <input type="checkbox"/> <b>At discontinuation</b>
Hepatitis A Ab	<input type="checkbox"/>			
Hepatitis B Panel <sup>2</sup>	<input type="checkbox"/>			<input type="checkbox"/> HBsAg w/ each pregnancy
Hepatitis C Ab <sup>3</sup>	<input type="checkbox"/>			<input type="checkbox"/> More frequent screening for PWIDs <input type="checkbox"/> Each pregnancy <input type="checkbox"/> Persistent ALT > 23IU/L

### Footnotes

1. Syphilis (reverse screening algorithm).  
Chlamydia and Gonorrhea at all applicable sites (genitourinary / oropharyngeal / rectal).
2. Surface antigen (HBsAg) + surface antibody (HBsAb) + core antibody (HBcAb)
3. If HCV-Ab is positive, obtain HCV-RNA.